

ON SYMPLECTIC MANIFOLDS WITH ASPHERICAL SYMPLECTIC FORM

YULI RUDYAK — ALEKSY TRALLE

Dedicated to the memory of Juliusz P. Schauder

ABSTRACT. We consider closed symplectically aspherical manifolds, i.e. closed symplectic manifolds (M, ω) satisfying the condition $[\omega]_{\pi_2 M} = 0$. Rudyak and Oprea [15] remarked that such manifolds have nice and controllable homotopy properties. Now it is clear that these properties are mostly determined by the fact that the strict category weight of $[\omega]$ equals 2. We apply the theory of strict category weight to the problem of estimating the number of closed orbits of charged particles in symplectic magnetic fields. In case of symplectically aspherical manifolds our theory enables us to improve some known estimations.

Introduction

A symplectically aspherical manifold is a symplectic manifold (M, ω) such that

$$\int_{S^2} f^* \omega = 0$$

for every map $f : S^2 \rightarrow M$. Such manifolds have explicitly appeared in papers of Floer [4] and Hofer [8] in context of Lagrangian intersection theory. These authors used an analytical advantage of symplectic asphericity: namely, it excludes the appearance of non-trivial pseudo-holomorphic spheres in M .

However, it turned out that symplectically aspherical manifolds have nice and controllable *homotopy* properties. Now it is clear that most of these properties

1991 *Mathematics Subject Classification*. Primary 55M30; Secondary 58F05.

Key words and phrases. Lusternik–Schnirelmann category, symplectic manifolds.

©1999 Juliusz Schauder Center for Nonlinear Studies

are based on the fact that strict category weight of $[\omega]$ equals 2 (Lemma 2.1), here $[\omega]$ is the de Rham cohomology class of the symplectic form ω . In [15] a weak version of this result (but with the same proof!) was used in order to prove the equality $\text{cat } M = \dim M$ for every closed symplectically aspherical manifold M . It was an important ingredient in proving of the Arnold conjecture about symplectic fixed points [12], [15].

In this paper we continue to study symplectically aspherical manifolds and, in particular, apply the theory of strict category weight to a special problem in symplectic topology. We show that additional topological arguments enable us to estimate the number of closed orbits of charged particles in *symplectic magnetic fields*. This subject has been recently of substantial interest [6], [9]. In a number of recent papers several estimates appeared and it was noted that these are probably not best possible. It follows from the results of this paper that at least in case of *symplectically aspherical* closed manifolds the corresponding estimates can indeed be sharpened (Theorem 3.4).

Notice that we don't prove any new analytic results, we only show that our topological observations do help to improve the existing estimates [9] in case of symplectically aspherical manifolds. We think that the technique of category weight potentially may have many applications to problems in symplectic topology, and, in a broader sense, to other non-linear analytic problems.

Acknowledgment. This article was written when both authors were visiting Max Planck Institut für Mathematik in Bonn. They are grateful to the Institute and to Professor Yuri Manin for the support of this work. Also, the research of the second author was partially financed by the Polish Research Committee (KBN).

We express our sincere thanks to Ely Kerman and Serge Tabachnikov for useful discussions.

1. Topological set-up

DEFINITION 1.1 ([10], [5]). Given a map $\varphi : A \rightarrow X$, we say that a subset U of A is φ -categorical if it is open (in A) and $\varphi|U$ is null-homotopic. We define the *Lusternik–Schnirelmann category* $\text{cat } \varphi$ of φ as follows:

$$\text{cat } \varphi = \min\{k \mid A = U_1 \cup \dots \cup U_{k+1} \text{ where each } U_i \text{ is } \varphi\text{-categorical}\}.$$

Furthermore, we define the *Lusternik–Schnirelmann category* $\text{cat } X$ of a space X by setting $\text{cat } X := \text{cat } 1_X$.

Given a path connected pointed space X , let

$$(1.2) \quad \varepsilon : S\Omega X \rightarrow X$$

be the map adjoint to $1_{\Omega X}$, where ΩX is the loop space of X and S denotes the (reduced) suspension, (see e.g. [18]).

THEOREM 1.3 ([17, Theorems 3, 19' and 21]). *Let $\varphi : A \rightarrow X$ be a map of connected Hausdorff paracompact spaces. Then $\text{cat } \varphi < 2$ iff there is a map $\psi : A \rightarrow S\Omega X$ such that $\varepsilon\psi \simeq \varphi$.*

DEFINITION 1.4 ([11]). Let X be a Hausdorff paracompact space, and let $u \in H^q(X; G)$ be an arbitrary element. We define the *strict category weight* of u (denoted by $\text{swgt } u$) by setting

$$\text{swgt } u = \sup\{k \mid \varphi^*u = 0 \text{ for every map } \varphi : A \rightarrow X \text{ with } \text{cat } \varphi < k\}$$

where A runs over all Hausdorff paracompact spaces.

We use the term “strict category weight”, since the term “category weight” is already used (introduced) by Fadell–Husseini [2]. Concerning the relation between category weight and strict category weight, see [11], [16].

We remark that $\text{swgt } u = \infty$ if $u = 0$.

THEOREM 1.5. *Let X be a path connected Hausdorff paracompact space, and let $u \in H^*(X)$ be an arbitrary cohomology class. Then the following holds:*

- (i) if $\text{deg } u > 0$ then $\text{swgt } u \geq 1$;
- (ii) for every map $f : Y \rightarrow X$ we have $\text{cat } f \geq \text{swgt } u$ provided that $f^*u \neq 0$;
- (iii) $\text{cat } X \geq \text{swgt } u$ provided that $u \neq 0$;
- (iv) for every map $f : Y \rightarrow X$ we have $\text{swgt } f^*u \geq \text{swgt } u$;
- (v) for every $v \in H^*(X)$ we have $\text{swgt}(uv) \geq \text{swgt } u + \text{swgt } v$;
- (vi) $\text{swgt } u \geq 2$ iff $\varepsilon^*u = 0$ where ε is as in (1.2).

PROOF. (i)–(iv) see [11], [14]. (vi) This follows from Theorem 1.3. \square

2. Strict category weight and symplectically aspherical manifolds

Given a space X and an element $u \in H^2(X; G)$, the notation $u|_{\pi_2(X)} = 0$ means that

$$\langle u, h(a) \rangle = 0 \quad \text{for every } a \in \pi_2(X)$$

where $h : \pi_2(X) \rightarrow H_2(X)$ is the Hurewicz homomorphism and $\langle -, - \rangle$ is the Kronecker pairing.

Notice that a symplectic manifold (M, ω) is symplectically aspherical if and only if $[\omega]|_{\pi_2(M)} = 0$.

LEMMA 2.1 (cf. [15]). *Let X be a finite CW-space, and let $u \in H^*(X; \mathbb{R})$ be a cohomology class such that $u|_{\pi_2(X)} = 0$. Then the following holds:*

- (i) *for every map $f : X \rightarrow K(\pi_1(X), 1)$ which induces an isomorphism of fundamental groups,*

$$u \in \text{Im}\{f^* \mid H^2(K(\pi_1(X); \mathbb{R})) \rightarrow H^2(X; \mathbb{R})\};$$

- (ii) $\text{swgt } u \geq 2$.

PROOF. (i) Let $K := K(\pi_1(X), 1)$, and let $f : X \rightarrow K$ be a map such that $f_* : \pi_1(X) \rightarrow \pi_1(K)$ is an isomorphism. Recall that for every space Y the evaluation homomorphism $e : H^*(Y; \mathbb{R}) \rightarrow \text{Hom}(H_*(Y), \mathbb{R})$ is an isomorphism. Furthermore, there is the Hopf exact sequence

$$\pi_2(X) \xrightarrow{h} H_2(X) \longrightarrow H_2(K) \longrightarrow 0.$$

So, we have the following commutative diagram with the exact row:

$$\begin{CD} H^2(K; \mathbb{R}) @>f^*>> H^2(X; \mathbb{R}) \\ @V\cong VV @VVe\cong V \\ \text{Hom}(H_2(K), \mathbb{R}) @>\text{Hom}(f_*, 1)>> \text{Hom}(H_2(X), \mathbb{R}) @>\text{Hom}(h, 1)>> \text{Hom}(\pi_2(X), \mathbb{R}). \end{CD}$$

Since $\text{Hom}(h, 1)(e^*u) = 0$, we conclude that $u \in \text{Im } f^*$.

- (ii) Consider the commutative diagram

$$\begin{CD} S\Omega X @>S\Omega f>> S\Omega K \\ @V\varepsilon_X VV @VV\varepsilon_K V \\ X @>f>> K \end{CD}$$

with $\varepsilon_X, \varepsilon_K$ as in (1.2). Because of (i), $u = f^*v$ for some $v \in H^2(K; \mathbb{R})$. Furthermore, ΩK is homotopy equivalent to the discrete space π , and so $S\Omega K$ is homotopy equivalent to a wedge of circles. Hence, $H^2(K; \mathbb{R}) = 0$, and, in particular, $\varepsilon_K^*v = 0$. Hence, $\varepsilon_X^*u = \varepsilon_X^*f^*v = (S\Omega f)^*(\varepsilon_K v) = 0$. Thus, by Theorem 1.5(vi), $\text{swgt } u \geq 2$. □

COROLLARY 2.2. *Let (M, ω) be a symplectic manifold, and let K denote the space $K(\pi_1(M), 1)$. The following three conditions are equivalent:*

- (i) *(M, ω) is symplectically aspherical;*
- (ii) *there exists a map $f : M \rightarrow K$ which induces an isomorphism of fundamental groups and such that*

$$[\omega] \in \text{Im}\{f^* : H^2(K; \mathbb{R}) \rightarrow H^2(M; \mathbb{R})\};$$

(iii) *there exists a map $f : M \rightarrow K$ such that*

$$[\omega] \in \text{Im}\{f^* : H^2(K; \mathbb{R}) \rightarrow H^2(M; \mathbb{R})\}$$

PROOF. (i) \Rightarrow (ii) follows from 2.1. (ii) \Rightarrow (iii) is trivial.

(iii) \Rightarrow (i) If $[\omega] = f^*u$ for some $u \in H^2(K; \mathbb{R})$ then $u|_{\pi_2(K)} = 0$ since $\pi_2(K) = 0$, and thus $[\omega]|_{\pi_2(M)} = 0$. □

THEOREM 2.3. *Let (M^{2m}, ω) be a closed symplectically aspherical manifold, and let*

$$\mathbb{C}P^n \xrightarrow{i} E \xrightarrow{p} M$$

be a $\mathbb{C}P^n$ -fibration over M . Then $\text{cat } E \geq 2m + n$.

PROOF. Let $u \in H^2(\mathbb{C}P^n; \mathbb{R})$ be a non-zero element. It is well-known that, for the fibration on hand, its Leray–Serre spectral sequence with real coefficients collapses, [19]. So, there is an element $v \in H^2(E; \mathbb{R})$ such that $i^*v = u$. Moreover, according to the Leray–Hirsch Theorem, see e.g. [18, (15.47)]

$$(p^*[\omega])^m v^n \neq 0 \in H^*(E; \mathbb{R}).$$

Now, by Lemma 2.1, $\text{swgt}[\omega] \geq 2$, and so, by Theorem 1.5(ii), (iv), (v),

$$\text{cat } E \geq \text{swgt}((p^*[\omega])^m v^n) \geq \text{swgt } p^*([\omega]^m) + \text{swgt } v^n \geq \text{swgt}[\omega]^m + n \geq 2m + n,$$

as claimed. □

3. Application to particle dynamics in magnetic field

In this section we show how one can apply the previous topological observations to estimation of number of closed trajectories of a charged particle in a symplectic magnetic field. Mathematically, this problem states as follows.

Let (M, ω) be a closed symplectic manifold, let $p : T^*M \rightarrow M$ be the cotangent bundle, and let φ be the standard 1-form on T^*M . Consider the symplectic form $\Omega := d\varphi + p^*\omega$ on T^*M . Given a Hamiltonian H on T^*M , consider the dynamics

$$(3.1) \quad i_X \Omega = dH.$$

The integral curves of the vector field X are the trajectories of a charged particle in a magnetic field with the Hamiltonian H .

Choose a Riemannian metric g and an almost complex structure J on M such that $\omega(Y, Z) = g(Y, JZ)$ for every vector fields Y, Z on M , and consider the so-called metric Hamiltonian $H : T^*M \rightarrow \mathbb{R}$, defined by $H(\xi) = \langle \xi, \xi \rangle_g$ for every covector ξ on M . Kerman [9] proved that, for every $\varepsilon > 0$ small enough, the energy level $H^{-1}(\varepsilon)$ contains at least $m + \text{cl}(M)$ closed trajectories of the vector field X as in (3.1); here cl denotes the cup-length. In greater detail, Kerman proved the following

THEOREM 3.2 ([9]). *For any metric Hamiltonian H and every ε small enough, there is a fiber bundle $p : \Sigma \rightarrow M$ with the following properties:*

- (1) $\Sigma \subset H^{-1}(\varepsilon)$;
- (2) $p : \Sigma \rightarrow M$ is a locally trivial bundle with the fiber S^{2m-1} ;
- (3) there is a free fiberwise S^1 -action on Σ , and the restriction of this action to any fiber coincides with the standard S^1 -action on S^{2m-1} ;
- (4) the energy level $H^{-1}(\varepsilon)$ contains at least $1 + \text{cat}(\Sigma/S^1)$ closed trajectories of the vector field X .¹

Notice that the projection $p : \Sigma \rightarrow M$ yields a locally trivial bundle

$$q : \Sigma/S^1 \rightarrow M$$

with the fiber $S^{2m-1}/S^1 = \mathbb{C}P^{m-1}$. Because of this, Kerman obtains the estimation

$$\text{cat}(\Sigma/S^1) \geq \text{cl}(\Sigma/S^1) \geq \text{cl} M + \text{cl}(\mathbb{C}P^{m-1}) = \text{cl} M + m - 1,$$

and, in particular, the following corollary.

COROLLARY 3.3. *The set $H^{-1}(\varepsilon)$ contains at least $m + \text{cl}(M)$ closed trajectories of X .*

We remark that this estimation can be improved if the symplectic form ω on M is aspherical.

THEOREM 3.4. *Let (M^{2m}, ω) be a closed symplectic manifold such that $[\omega]|_{\pi_2(M)} = 0$, and let H be a metric Hamiltonian on T^*M . Then, for every ε small enough, the set $H^{-1}(\varepsilon)$ contains at least $3m$ closed trajectories of the vector field X as in (3.1).*

PROOF. Since Σ/S^1 is fibered over M with the fiber $\mathbb{C}P^{m-1}$, we conclude that, by Theorem 2.3, $\text{cat}(\Sigma/S^1) \geq 3m - 1$. Now the result follows from Theorem 3.2(4). □

REMARKS 3.5. (1) Using the language of mathematical physics, Theorem 3.4 can be reformulated as follows. *A charged particle on a symplectically aspherical manifold, under the influence of the symplectic magnetic field, has at least $3m$ closed trajectories on any sufficiently low kinetic energy level.*

(2) Clearly, $\text{cl}(M) \leq 2m$ for every connected $2m$ -dimensional manifold M . Furthermore, there are many examples of closed symplectic manifolds with aspherical symplectic form and such that $\text{cl} M < 2m$, see Proposition 4.3 below. So, for closed symplectically aspherical manifolds the estimation from Theorem 3.4 is stronger than this one from Corollary 3.3.

¹Notice that the Lusternik–Schnirelmann category considered in our paper is one less than the Lusternik–Schnirelmann category used in [9].

4. Examples

In this section we present examples of symplectically aspherical closed manifolds. These examples give the reader a picture of the class of manifolds covered by the results of the present paper.

To start with, notice that there are two constructions which preserve symplectic asphericity. First, the Cartesian product of two symplectically aspherical manifolds is symplectically aspherical. Second, if $f : (M, \omega) \rightarrow (N, \sigma)$ is a map of symplectic manifolds with $f^*\sigma = \omega$, then (M, ω) is symplectically aspherical provided (N, σ) is. This can be proved directly or deduced from Corollary 2.2.

As a special case, consider a symplectically aspherical manifold (M, ω) and a branched covering $p : \widetilde{M} \rightarrow M$ such that the branch locus is a symplectic submanifold of codimension 2. Then \widetilde{M} possesses a symplectic form $\widetilde{\omega}$ with $[\widetilde{\omega}] = p^*[\omega]$, [7]. Thus, $(\widetilde{M}, \widetilde{\omega})$ is symplectically aspherical.

Notice that if a manifold, say, M is covered by \mathbb{R}^n , then it is necessarily aspherical, i.e. $\pi_i M = 0$ for $i > 1$. Therefore, for any symplectic manifold covered by \mathbb{R}^n we have $\pi_2 M = 0$ and, in particular, the condition $[\omega]|_{\pi_2 M} = 0$ is trivially satisfied. So, taking quotients of simply connected solvable Lie groups by lattices (if there are any), we get many examples of aspherical closed manifolds.

EXAMPLE 4.1 (nilmanifolds and solvmanifolds). Let N be any simply connected nilpotent Lie group with the Lie algebra \mathfrak{n} satisfying the following condition: there exists a basis e_1, \dots, e_n of \mathfrak{n} such that all the structural constants c_{ij}^k of \mathfrak{n} with respect to this basis are *rational numbers*. By the Malcev Theorem, N necessarily admits a lattice, i.e. a discrete co-compact subgroup Γ . Hence we can form a closed aspherical nilmanifold N/Γ . One can easily get a characterization of *symplecticness* of such manifolds. It is known that each cohomology class in $H^*(N/\Gamma)$ contains a *homogeneous* representative, i.e. a differential form whose pullback to N is an invariant differential form on Lie group N , see [20, Chapter 2, Theorem 1.3]. Hence, every cohomologically symplectic manifold of the form N/Γ is symplectic. (Recall that a closed connected smooth manifold M^{2n} is called *cohomologically symplectic* if it possesses a 2-form ω such that $[\omega]^n \neq 0$.) One can check the cohomological symplecticness by calculating the Chevalley–Eilenberg complex for \mathfrak{n} . So, we have the following fact.

PROPOSITION 4.2. *Any closed cohomologically symplectic nilmanifold $M = N/\Gamma$ is symplectically aspherical.*

Now we present an explicit example of a symplectically aspherical closed 4-dimensional manifold K for which $\text{cl} K = 3$ and $\text{cat} K = 4$. Consider the Kodaira–Thurston manifold $K = (N_3/\Gamma) \times S^1$ defined as follows. Let N_3 stand

for the 3-dimensional Heisenberg group of all matrices of the form

$$\begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}$$

where $a, b, c \in \mathbb{R}$ and Γ denotes the subgroup of N_3 consisting of all matrices with integer entries. The cohomology of K is easy to calculate as was indicated below (see also [20]): the Chevalley–Eilenberg complex of K is given by the formula

$$\Lambda(\alpha, \beta, \gamma, \delta), \quad d\alpha = d\beta = d\delta = 0, \quad d\gamma = \alpha \wedge \beta,$$

where Λ denotes the exterior algebra generated by $\alpha, \beta, \gamma, \delta$ and these generators have degree 1. Hence the vector space $H^1(K; \mathbb{R}) \simeq \mathbb{R}^3$ is generated by $[\alpha], [\beta], [\delta]$ and, obviously, the length of the longest product in $H^*(K; \mathbb{R})$ is ≤ 3 . In fact, $\text{cl}(K) = 3$, the element $[\alpha][\delta][\beta\gamma]$ gives us the product of length 3. Notice that K is symplectic, since the cohomology class $[\alpha \wedge \delta + \beta \wedge \gamma]$ is symplectic. However, $\text{cat}(K) = 4$, since $\text{cat } V = \dim V$ for any closed aspherical manifold V (see [1]). Thus, $3 = \text{cl } K < \text{cat } K = 4$.

The previous example can be obviously generalized to any symplectic nilmanifold. Indeed, the Chevalley–Eilenberg complex of any nilmanifold N/Γ has the form $(\Lambda(x_1, \dots, x_n), d)$ with $n = \dim(N/\Gamma) = \dim N$ and the degrees of generators x_i being 1. If N/Γ is not a torus (i.e., if Γ is not Abelian), then $\text{Ker } d \neq 0$, and so $dx_i \neq 0$ for at least one x_i . Hence, any cup-product of any n cocycles is a boundary, and we obtain the following proposition.

PROPOSITION 4.3. *For any cohomologically symplectic non-toral nilmanifold N/Γ*

$$\text{cl}(N/\Gamma) < n = \dim(N/\Gamma) = \text{cat}(N/\Gamma).$$

In the same fashion one can analyze *solvmanifolds*. Here, however, it is not so easy to prove the existence of lattices and symplectic structures (cf. [20]). Nevertheless, there is a certain class of solvmanifolds where the answer is essentially the same as that for nilmanifolds. We say that a simply connected solvable Lie group G is *completely solvable*, if for the Lie algebra \mathfrak{g} of G all the operators

$$\text{ad } V : \mathfrak{g} \rightarrow \mathfrak{g}, \quad V \in \mathfrak{g},$$

have only real eigenvalues. Now, the following proposition can be proved similarly to Proposition 4.2.

PROPOSITION 4.4. *Let Γ be a co-compact discrete subgroup of a completely solvable Lie group G . If the manifold $M = G/\Gamma$ is cohomologically symplectic, then it is symplectically aspherical.*

Notice that there are also examples of symplectically aspherical closed manifolds with non-trivial π_2 , but the corresponding explicit constructions are rather

delicate and have very recently appeared [7]. The general description of the whole class of symplectically aspherical manifolds is given in Corollary 2.2. However, it seems rather difficult to use it to construct explicit examples. Recently Gompf produced such examples as branched coverings.

THEOREM 4.5 ([7]). *Let $p : \tilde{X} \rightarrow X$ be a d -fold branched covering of an orientable 4-manifold X such that the branch locus $B \subset X$ is obtained from a generically immersed surface $B^* \subset X$ by smoothing all double points. If B^* has $k \geq 1$ double points then $\pi_2(\tilde{X}) \otimes \mathbb{R} \neq 0$.*

So, if one takes any 4-dimensional closed symplectically aspherical manifold (X, ω) (e.g. the Kodaira–Thurston manifold) and a branched covering $p : \tilde{X} \rightarrow X$ as in Theorem 4.5, then $(\tilde{X}, \tilde{\omega})$ (described in the beginning of the section) is a symplectically aspherical manifold with $\pi_2(\tilde{X}) \neq 0$.

EXAMPLE 4.6. We can get more aspherical closed manifolds by considering lattices in *semisimple* Lie groups of non-compact type. It is known [3] that any simply connected semisimple Lie group contains a torsion-free, discrete, co-compact subgroup. Consequently, any symmetric space of non-compact type is the universal covering space of a locally symmetric space M . Hence, M is a closed and aspherical manifold.

In the same fashion, we can consider *hyperbolic manifolds* (examples: Riemannian surfaces).

REFERENCES

- [1] S. EILENBERG AND T. GANEVA, *On the Lusternik–Schnirelmann category of abstract groups*, Ann. of Math. **65** (1957), 517–518.
- [2] E. FADELL AND S. HUSSEINI, *Category weight and Steenrod operations*, Bol. Soc. Mat. Mexicana **37** (1992), 151–161.
- [3] F. T. FARRELL AND L. E. JONES, *Classical Aspherical Manifolds*, Amer. Math. Soc., 1990.
- [4] A. FLOER, *Symplectic fixed points and holomorphic spheres*, Comm. Math. Phys. **120** (1989), 575–611.
- [5] R. FOX, *On the Lusternik–Schnirelmann category*, Ann. of Math. **42** (1941), 333–370.
- [6] V. GINZBURG AND E. KERMAN, *Periodic orbits in magnetic fields in dimensions greater than two*, Preprint, 1999, Math. DG/9902148.
- [7] R. GOMPF, *On symplectically aspherical manifolds with nontrivial π_2* , Math. Res. Lett. **5** (1999), 599–603.
- [8] H. HOFER, *Lusternik–Schnirelmann theory for Lagrangian intersections*, Ann. Inst. H. Poincaré–Anal. Non Linéaire **5** (1988), 465–499.
- [9] E. KERMAN, *Periodic orbits of Hamiltonian flows near symplectic critical submanifolds*, Preprint, 1999, Math. DG/9903100.
- [10] L. A. LUSTERNIK AND L. G. SCHNIRELMANN, *Methodes Topologiques dans le Problèmes Variationels*, Hermann, Paris, 1934.

- [11] YU. RUDYAK, *On category weight and its applications*, *Topology* **38** (1999), 37–55.
- [12] ———, *On analytical applications of stable homotopy (the Arnold conjecture, critical points)*, *Math. Z.* **230** (1999), 659–672.
- [13] ———, *Category weight: new ideas concerning Lusternik–Schnirelmann category*, *Homotopy and Geometry*, Banach Center Publications, vol. 45, Warszawa, 1998, pp. 47–61.
- [14] ———, *On strict category weight, gradient like flows and the Arnold conjecture*, *Internat. Math. Res. Notices* (to appear).
- [15] YU. RUDYAK AND J. OPREA, *On the Lusternik–Schnirelmann category of symplectic manifolds and the Arnold conjecture*, *Math. Z.* **230** (1999), 673–678.
- [16] J. STROM, *Category Weight and Essential Category Weight*, Thesis, Univ. of Madison–Wisconsin, 1997.
- [17] A. ŠVARC, *The genus of a fiber space*, *Amer. Math. Soc. Transl. Ser.* **55** (1966), 49–140.
- [18] R. SWITZER, *Algebraic Topology — Homotopy and Homology*, Springer-Verlag, Berlin, 1975.
- [19] J.-C. THOMAS, *Rational homotopy of Serre fibrations*, *Ann. Inst. Fourier (Grenoble)* **31** (1981), 71–90.
- [20] A. TRALLE AND J. OPREA, *Symplectic Manifolds with no Kähler Structure*, *Lecture Notes in Math.* **1661**, Springer-Verlag, Berlin, 1997.

Manuscript received October 15, 1999

YULI RUDYAK
FB6/Mathematik
Universität Siegen
57068 Siegen, GERMANY

E-mail address: rudyak@mathematik.uni-siegen.de, july@mathi.uni-heidelberg.de

ALEKSY TRALLE
University of Warmia and Masuria
10-561 Olsztyn, POLAND

E-mail address: tralle@tufi.wsp.olsztyn.pl, tralle@matman.wsp.olsztyn.pl