Topological Methods in Nonlinear Analysis Journal of the Juliusz Schauder Center Volume 15, 2000, 401–401

CORRECTIONS TO "STABLE MAPS OF GENUS ZERO TO FLAG SPACES" (TOPOL. METHODS NONLINEAR ANAL. 11 (1998), 207–217)

YURI I. MANIN

- (1) Divide the left hand side of (2.3) by [PGL(2)]. The same in the unnumbered formula several lines after (2.11) preceding the sentence "which is (2.3)".
- (2) Divide by [W] the left hand side of the unnumbered formula following (2.11). Divide by $P_W(q)$ the left hand side of (2.13) and (2.17).
- (3) Divide the left hand side of (2.12) by $q^2(q^4-1)$.
- (4) Replace (2.15) by

$$(1 + q^2 t - q^2 \psi)\psi_t = 1 + \psi.$$

(5) Replace (2.16) by

(2.15)

(2.16)
$$(1+t+\varphi^0)\log(1+t+\varphi^0) = 2\varphi^0 + t - X(1+t+\varphi^0),$$
$$X = \frac{1}{q^2-1} \left(\frac{E(W,z)}{P_W(q)} - 1\right)\Big|_{q^2=1}.$$

I thank Nikita Nekrasov who altered me to these mistakes.

Manuscript received January 18, 2000

YURI I. MANIN Max-Planck-Institut für Mathemtik Bonn, GERMANY

 $E\text{-mail}\ address:$ manin@mpim-bonn.mpg.deTMNA: Volume 15 – 2000 – N° 2

O2000Juliusz Schauder Center for Nonlinear Studies

401