

**DOUBLE POSITIVE SOLUTIONS  
FOR SECOND ORDER NONLOCAL FUNCTIONAL  
AND ORDINARY BOUNDARY VALUE PROBLEMS**

PANAGIOTIS CH. TSAMATOS

---

ABSTRACT. In this paper we prove the existence of two positive solutions for a second order nonlinear functional nonlocal boundary value problem. The results are obtained by using a fixed point theorem on a Banach space, ordered by an appropriate cone, due to Avery and Henderson [1]. Using this theorem we have the advantage that the obtained two solutions have their values at three points of their domain upper and lower bounded by a-priori given constants.

**1. Introduction**

Let  $\mathbb{R}$  be the set of real numbers,  $\mathbb{R}^+ =: [0, \infty)$  and  $I =: [0, 1]$ . Also, let  $q \in [0, 1)$  and  $J := [-q, 0]$ . For every closed interval  $B \subseteq J \cup I$  we denote by  $C(B)$  the Banach space of all continuous real functions  $\psi: B \rightarrow \mathbb{R}$  endowed with the usual sup-norm

$$\|\psi\|_B := \sup\{|\psi(s)| : s \in B\}.$$

Also, we define the set  $C^+(B)$  as follows

$$C^+(B) := \{\psi \in C(B) : \psi \geq 0\}.$$

---

2000 *Mathematics Subject Classification.* 34K10, 34B18.

*Key words and phrases.* Positive solutions, boundary value problems, second order functional differential equations, nonlocal boundary conditions.

If  $x \in C(J \cup I)$  and  $t \in I$ , then we denote by  $x_t$  the element of  $C(J)$  defined by

$$x_t(s) = x(t + s) \quad \text{for } s \in J.$$

Now, consider the equation

$$(1.1) \quad x''(t) + f(t, x_t) = 0 \quad \text{for } t \in I,$$

along with the initial condition

$$(1.2) \quad x_0 = \phi,$$

and the nonlocal boundary condition

$$(1.3) \quad x'(1) = \int_0^1 x'(s) dg(s),$$

where  $f: \mathbb{R}^+ \times C^+(J) \rightarrow \mathbb{R}^+$  and  $\phi: J \rightarrow \mathbb{R}^+$  are continuous functions,  $g: I \rightarrow \mathbb{R}$  is a nondecreasing function, such that  $g(0) = 0$  and  $1 - g(1) > 0$ .

The problem of existence of positive solutions for boundary value problems for second order differential equations which involve a nonlocal condition like (1.3) has been treated recently by Karakostas and Tsamatos [8]–[11] and Tsamatos [20]. Moreover, boundary value problems with integral boundary conditions for second order differential equations with retarded arguments are the subject of the papers [12] and [17]. In the recent years an increasing interest is also observed for boundary value problems concerning functional differential equations (see [6], [7] and the references therein). Fixed point theorems on Banach spaces ordered by appropriate cones are usually the tools for proving multiple positive solutions for boundary value problems. The famous Guo–Krasnoselskii fixed point theorem [13], [22] seems to be used in the majority of the papers on this subject. Also the well known Leggett–Williams fixed point theorem [14] and some recent generalizations of it are used in proving existence of multiple positive solutions for various types of boundary value problems.

For a detailed exposition of the theory of functional differential equations, like (1.1), the reader is referred to the books due to Hale and Lunel [4] and Azbelev *et al.* [3].

In this paper, we choose to use a fixed point theorem, on a Banach space ordered by cones, due to Avery and Henderson [1] (see also [15], [19]) which, apart from guaranteeing the existence of two positive solutions, has the advantage to offer some additional information on these solutions. In our results, the values of these solutions at three given points of their domain are upper or lower bounded by a-priori given constants. We note that this fixed point theorem was used recently in several papers (see [2], [5], [15], [16], [18], [19] and the references therein).

The present paper is motivated mainly by the papers [8]–[11] in which the problem of the existence of multiple positive solutions for ordinary differential equations with the nonlocal boundary condition (1.3) is studied.

Since the results we present are new even in the ordinary differential equations case, we mention them for this case too, underlining the necessary adjustments that have to be made to the hypothesis referring to the functional case.

The paper is organized as follows. In Section 2 we present the definitions and the lemmas we are going to use, as well as the fixed point theorem, on which we base our results. In Section 3, we present the new results for the functional case and then in Section 4 the results for the ordinary case. Finally, in Section 5 we give some applications of our results.

## 2. Preliminaries and some basic lemmas

**DEFINITION 2.1.** A function  $x \in C(J \cup I)$  is a solution of the boundary value problem (1.1)–(1.3) if  $x$  satisfies equation (1.1), the boundary condition (1.3) and, moreover  $x|_J = \phi$ .

**LEMMA 2.2.** A function  $x \in C(J \cup I)$  is a solution of the boundary value problem (1.1)–(1.3) if and only if  $x$  is a fixed point of the operator  $A: C(J \cup I) \rightarrow C(J \cup I)$ , with

$$Ax(t) = \begin{cases} \phi(t) & \text{for } t \in J, \\ \phi(0) + \zeta t \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_0^t \int_s^1 f(r, x_r) dr ds & \text{for } t \in I, \end{cases}$$

where  $\zeta := 1/(1 - g(1))$ .

**PROOF.** Suppose that  $x$  is a solution of the boundary value problem (1.1)–(1.3). Then, obviously,  $x|_J = \phi$ . Moreover, by integrating (1.1) we get

$$(2.1) \quad x'(t) = x'(1) + \int_t^1 f(s, x_s) ds \quad \text{for } t \in I.$$

Also from (1.3) and (2.1) we have

$$x'(1) = \int_0^1 x'(s) dg(s) = x'(1) \int_0^1 dg(s) + \int_0^1 \int_s^1 f(r, x_r) dr dg(s).$$

Therefore

$$(2.2) \quad x'(1) = \zeta \int_0^1 \int_s^1 f(r, x_r) dr dg(s).$$

Combining (2.1) and (2.2) we conclude that

$$x'(t) = \zeta \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_t^1 f(s, x_s) ds \quad \text{for } t \in I,$$

which by integration from 0 to  $t$ ,  $t \in I$  gives

$$x(t) = \phi(0) + \zeta t \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_0^t \int_s^1 f(r, x_r) dr ds.$$

The above step gives that, if  $x$  is a solution of the boundary value problem (1.1)–(1.3), then  $x = Ax$ .

Reciprocally, suppose that  $x$  is a fixed point of the operator  $A$ . Then, obviously,  $\phi(t) = x(t) = x(0+t) = x_0(t)$  for  $t \in J$ . Also from the form of  $A$  we have

$$(2.3) \quad x'(t) = \zeta \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_t^1 f(r, x_r) dr \quad \text{for } t \in I.$$

Therefore

$$(2.4) \quad x'(1) = \zeta \int_0^1 \int_s^1 f(r, x_r) dr dg(s).$$

Then from (2.3), (2.4) and the fact that  $\zeta := 1/(1-g(1))$ ,  $g(0) = 0$ , we get

$$(2.5) \quad x'(t) = x'(1) + \int_t^1 f(s, x_s) ds \quad \text{for } t \in I.$$

Using (2.4) and (2.5) we can easily obtain

$$x'(1) = \int_0^1 x'(s) dg(s).$$

Finally, from (2.5) we have

$$x'(t) = x'(1) - \int_1^t f(s, x_s) ds \quad \text{for } t \in I$$

and so

$$x''(t) + f(t, x_t) = 0 \quad \text{for } t \in I.$$

The proof is complete. □

The following lemma can be found in [21].

LEMMA 2.3. *If a function  $x \in C(I)$  is concave, nondecreasing and nonnegative then*

$$x(t) \geq t\|x\|, \quad 0 \leq t \leq 1.$$

The results proved in this paper are based on the following Theorem 2.7 due to R. I. Avery and J. Henderson [1] (see also [15] and [19]). As we mentioned in the introduction, this theorem ensures that our boundary value problem (1.1)–(1.3) has at least two distinct positive solutions and, moreover, for each of these solutions, we have an upper bound at some specific point of its domain and a lower bound at some other specific point of its domain. Also, both solutions are concave and nondecreasing on  $I$ . In order to apply this theorem some definitions are necessary.

DEFINITION 2.4. Let  $\mathbb{E}$  be a real Banach space. A cone in  $\mathbb{E}$  is a nonempty, closed set  $\mathbb{P} \subset \mathbb{E}$  such that

- (a)  $\kappa u + \lambda v \in \mathbb{P}$  for all  $u, v \in \mathbb{P}$  and all  $\kappa, \lambda \geq 0$ ,
- (b)  $u, -u \in \mathbb{P}$  implies  $u = 0$ .

DEFINITION 2.5. Let  $\mathbb{P}$  be a cone in a real Banach space  $\mathbb{B}$ . A functional  $\psi: \mathbb{P} \rightarrow \mathbb{B}$  is said to be increasing on  $\mathbb{P}$  if  $\psi(x) \leq \psi(y)$ , for any  $x, y \in \mathbb{P}$  with  $x \leq y$ , where  $\leq$  is the partial ordering induced to the Banach space by the cone  $\mathbb{P}$ , i.e.

$$x \leq y \quad \text{if and only if} \quad y - x \in \mathbb{P}.$$

DEFINITION 2.6. Let  $\psi$  be a nonnegative functional on a cone  $\mathbb{P}$ . For each  $d > 0$  we denote by  $\mathbb{P}(\psi, d)$  the set

$$\mathbb{P}(\psi, d) := \{x \in \mathbb{P} : \psi(x) < d\}.$$

THEOREM 2.7. Let  $\mathbb{P}$  be a cone in a real Banach space  $\mathbb{E}$ . Let  $\alpha$  and  $\gamma$  be increasing, nonnegative, continuous functionals on  $\mathbb{P}$ , and let  $\theta$  be a nonnegative functional on  $\mathbb{P}$  with  $\theta(0) = 0$  such that, for some  $c > 0$  and  $\Theta > 0$ ,

$$\gamma(x) \leq \theta(x) \leq \alpha(x) \quad \text{and} \quad \|x\| \leq \Theta\gamma(x),$$

for all  $x \in \overline{\mathbb{P}(\gamma, c)}$ . Suppose there exists a completely continuous operator

$$A: \overline{\mathbb{P}(\gamma, c)} \rightarrow \mathbb{P}$$

and real constants  $a, b$ , with  $0 < a < b < c$ , such that

$$\theta(\lambda x) \leq \lambda\theta(x) \quad \text{for } 0 \leq \lambda \leq 1 \text{ and } x \in \partial\mathbb{P}(\theta, b),$$

and

- (a)  $\gamma(Ax) > c$  for all  $x \in \partial\mathbb{P}(\gamma, c)$ ,
- (b)  $\theta(Ax) < b$  for all  $x \in \partial\mathbb{P}(\theta, b)$ ,
- (c)  $\mathbb{P}(\alpha, a) \neq \emptyset$  and  $\alpha(Ax) > a$  for all  $x \in \partial\mathbb{P}(\alpha, a)$ ,

or

- (a')  $\gamma(Ax) < c$  for all  $x \in \partial\mathbb{P}(\gamma, c)$ ,
- (b')  $\theta(Ax) > b$  for all  $x \in \partial\mathbb{P}(\theta, b)$ ,
- (c')  $\mathbb{P}(\alpha, a) \neq \emptyset$ , and  $\alpha(Ax) < a$  for all  $x \in \partial\mathbb{P}(\alpha, a)$ .

Then  $A$  has at least two fixed points  $x_1$  and  $x_2$  belonging to  $\overline{\mathbb{P}(\gamma, c)}$  such that

$$a < \alpha(x_1), \quad \theta(x_1) < b < \theta(x_2) \quad \text{and} \quad \gamma(x_2) < c.$$

### 3. Main results

Define the set  $\mathbb{K} := \{x \in C(J \cup I) : x(t) \geq 0, t \in J \cup I, x|_I \text{ is concave and nondecreasing}\}$ , which is a cone in  $C(J \cup I)$ . Also let  $0 < r_1 \leq r_2 \leq r_3 \leq 1$  and consider the following functionals

$$\begin{aligned}\gamma(x) &= x(r_1) \quad \text{for } x \in \mathbb{K}, \\ \theta(x) &= x(r_2) \quad \text{for } x \in \mathbb{K}, \\ \alpha(x) &= x(r_3) \quad \text{for } x \in \mathbb{K}.\end{aligned}$$

It is easy to see that  $\alpha, \gamma$  are nonnegative, increasing and continuous functionals on  $\mathbb{K}$ ,  $\theta$  is nonnegative on  $\mathbb{K}$  and  $\theta(0) = 0$ . Also, it is straightforward that

$$(3.1) \quad \gamma(x) \leq \theta(x) \leq \alpha(x),$$

since  $x \in \mathbb{K}$  is nondecreasing on  $I$ . Furthermore, for any  $x \in \mathbb{K}$ , by Lemma 2.3, we have

$$\gamma(x) = x(r_1) \geq r_1 \|x\|_I.$$

So

$$(3.2) \quad \|x\|_I \leq \frac{1}{r_1} \gamma(x) \quad \text{for } x \in \mathbb{K}.$$

Additionally, by the definition of  $\theta$  it is obvious that

$$\theta(\lambda x) = \lambda \theta(x), \quad 0 \leq \lambda \leq 1, \quad x \in \mathbb{K}.$$

Now, if  $D \subset I$ , consider the functions  $H: C(I) \rightarrow C(I)$  and  $H_D: C(I) \rightarrow C(I)$  by

$$(Hz)(s) := \int_s^1 z(r) dr \quad \text{for } s \in I$$

and

$$(H_D z)(s) := \int_{D \cap [s, 1]} z(r) dr \quad \text{for } s \in I.$$

At this point, we state the following assumptions:

- (H<sub>1</sub>) There exist  $M > 0$ , a continuous function  $u: I \rightarrow \mathbb{R}^+$  and a nondecreasing function  $L: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \leq u(t)L(\|y\|_J) \quad \text{for } t \in I, \quad y \in C^+(J)$$

and also

$$\phi(0) + L(M) \left( \zeta r_2 \int_0^1 (Hu)(s) dg(s) + \int_0^{r_2} (Hu)(s) ds \right) < Mr_2.$$

- (H<sub>2</sub>) There exist a constant  $\delta \in (0, 1)$  and functions  $\tau: I \rightarrow [0, q]$ , continuous  $v: I \rightarrow \mathbb{R}^+$  and nondecreasing  $w: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \geq v(t)w(y(-\tau(t))), \quad \text{for } t \in X, \quad y \in \{h \in C^+(J) : \|h\|_J < M\},$$

where  $X := \{t \in I : \delta \leq t - \tau(t) \leq 1\}$  and  $M$  is defined in  $(H_1)$ .

$(H_3)$  There exist  $\rho_1, \rho_3 > 0$  such that

$$\frac{\rho_i}{\delta} < \phi(0) + w(\rho_i) \left( r_i \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_i} (H_X v)(s) ds \right), \quad i = 1, 3,$$

and  $\rho_3/\delta < Mr_2 < \rho_1/\delta$ .

Notice that if  $\phi(0) \neq 0$ , then these  $\rho_1, \rho_3$  always exist.

REMARK 3.1. *It is easy to see that  $\sup\{v(t) : t \in X\} > 0$  and  $\text{meas}(X \cap [s, 1]) > 0, s \in [0, 1)$  in the assumption  $(H_2)$ , imply that*

$$\int_0^1 (H_X v)(s) dg(s) > 0 \quad \text{and} \quad \int_0^{r_i} (H_X v)(s) ds > 0, \quad i = 1, 3.$$

THEOREM 3.2. *Suppose that assumptions  $(H_1)$ – $(H_3)$  hold and furthermore  $\|\phi\|_J \leq M$ . Then the boundary value problem (1.1)–(1.3) has at least two concave and nondecreasing on  $I$  and positive on  $J \cup I$  solutions  $x_1, x_2$  such that  $x_1(r_3) > \rho_3/\delta, x_1(r_2) < Mr_2, x_2(r_2) > Mr_2$  and  $x_2(r_1) < \rho_1/\delta$ .*

PROOF. First of all, we observe that, because of  $(H_1)$ ,  $f(t, \cdot)$  maps bounded sets into bounded sets. Therefore  $A$  is a completely continuous operator.

Now we set  $a = \rho_3/\delta, b = Mr_2, c = \rho_1/\delta$  and we consider a  $x \in \overline{\mathbb{K}(\gamma, c)}$ . Then since  $\zeta > 0$  and  $f(t, x_t) \geq 0$  for every  $t \in I$ , we get that  $Ax(t) \geq 0, t \in I$ . Also  $Ax(t) = \phi(t) \geq 0, t \in J$ . Thus  $Ax(t) \geq 0, t \in J \cup I$ . Moreover,  $(Ax)''(t) = -f(t, x_t) \leq 0$ , which means that  $Ax$  is concave on  $I$ . Also it is clear that  $(Ax)'(t) \geq 0$  for  $t \in I$ . So  $A: \overline{\mathbb{K}(\gamma, c)} \rightarrow \mathbb{K}$ .

Now let  $x \in \partial\mathbb{K}(\gamma, c)$ . Then  $\gamma(x) = x(r_1) = c$  and so  $\|x\|_I \geq c$ . Having in mind assumption  $(H_2)$ , we get

$$\begin{aligned} \gamma(Ax) &= Ax(r_1) = \phi(0) + \zeta r_1 \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_0^{r_1} \int_s^1 f(r, x_r) dr ds \\ &\geq \phi(0) + \zeta r_1 \int_0^1 \int_{X \cap [s, 1]} f(r, x_r) dr dg(s) + \int_0^{r_1} \int_{X \cap [s, 1]} f(r, x_r) dr ds \\ &\geq \phi(0) + \zeta r_1 \int_0^1 \int_{X \cap [s, 1]} v(r)w(x_r(-\tau(r))) dr dg(s) \\ &\quad + \int_0^{r_1} \int_{X \cap [s, 1]} v(r)w(x_r(-\tau(r))) dr ds \\ &= \phi(0) + \zeta r_1 \int_0^1 \int_{X \cap [s, 1]} v(r)w(x(r - \tau(r))) dr dg(s) \\ &\quad + \int_0^{r_1} \int_{X \cap [s, 1]} v(r)w(x(r - \tau(r))) dr ds \\ &\geq \phi(0) + \zeta r_1 \int_0^1 \int_{X \cap [s, 1]} v(r)w(x(\delta)) dr dg(s) \end{aligned}$$

$$+ \int_0^{r_1} \int_{X \cap [s,1]} v(r)w(x(\delta)) dr ds.$$

Additionally, by assumption (H<sub>3</sub>) and Lemma 2.3, we have

$$\begin{aligned} \gamma(Ax) &\geq \phi(0) + w(\delta\|x\|_I) \left( r_1 \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_1} (H_X v)(s) ds \right) \\ &\geq \phi(0) + w(\delta c) \left( r_1 \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_1} (H_X v)(s) ds \right) \\ &= \phi(0) + w(\rho_1) \left( r_1 \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_1} (H_X v)(s) ds \right) > \frac{\rho_1}{\delta} = c. \end{aligned}$$

This means that condition (a) of Theorem 2.7 is satisfied.

Now let  $x \in \partial\mathbb{K}(\theta, b)$ . Then  $\theta(x) = x(r_2) = b$  and so by Lemma 2.3 we get

$$\|x\|_I \leq \frac{1}{r_2} x(r_2) = \frac{1}{r_2} \theta(x) = \frac{b}{r_2}.$$

Also we assumed that  $\|\phi\|_J \leq M = b/r_2$ , so  $\|x\|_{J \cup I} \leq b/r_2$ . Now, by (H<sub>1</sub>), we have

$$\begin{aligned} \theta(Ax) &= Ax(r_2) \\ &= \phi(0) + \zeta r_2 \int_0^1 \int_s^1 f(r, x_r) dr dg(s) + \int_0^{r_2} \int_s^1 f(r, x_r) dr ds \\ &\leq \phi(0) + \zeta r_2 \int_0^1 \int_s^1 u(r) L(\|x_r\|_J) dr dg(s) + \int_0^{r_2} \int_s^1 u(r) L(\|x_r\|_J) dr ds \\ &\leq \phi(0) + \zeta r_2 \int_0^1 \int_s^1 u(r) L\left(\frac{b}{r_2}\right) dr dg(s) + \int_0^{r_2} \int_s^1 u(r) L\left(\frac{b}{r_2}\right) dr ds \\ &= \phi(0) + L(M) \left( \zeta r_2 \int_0^1 (Hu)(s) dg(s) + \int_0^{r_2} (Hu)(s) ds \right) < Mr_2 = b. \end{aligned}$$

So condition (b) of Theorem 2.7 is also satisfied.

Now, define the function  $y: J \cup I \rightarrow \mathbb{R}$  with  $y(t) = a/2$ . Then it is obvious that  $\alpha(y) = a/2 < a$ , so  $\mathbb{K}(\alpha, a) \neq \emptyset$ . Also, for any  $x \in \partial\mathbb{K}(\alpha, a)$  we have  $\alpha(x) = x(r_3) = a$ . Therefore  $\|x\|_I \geq a$ . Now, having in mind assumption (H<sub>2</sub>) and as in the case of the functional  $\gamma$  above, we get

$$\begin{aligned} \alpha(Ax) = Ax(r_3) &\geq \phi(0) + \zeta r_3 \int_0^1 \int_{X \cap [s,1]} v(r)w(x(\delta)) dr dg(s) \\ &\quad + \int_0^{r_3} \int_{X \cap [s,1]} v(r)w(x(\delta)) dr ds. \end{aligned}$$

Then, by assumption (H<sub>3</sub>) and inequality of Lemma 2.3, we also have

$$\alpha(Ax) \geq \phi(0) + w(\delta a) \left( r_3 \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_3} (H_X v)(s) ds \right) > \frac{\rho_3}{\delta} = a.$$

Consequently, assumption (c) of Theorem 2.7 is satisfied.



The result can now be obtained by applying Theorem 2.7. □

The above Theorem 3.2 has been obtained by using the requirements (a)–(c) of Theorem 2.7. Using the requirements (a′)–(c′) of the same theorem we can also obtain another existence theorem (Theorem 3.3 below) for our boundary value problem (1.1)–(1.3). For this purpose we need the following assumptions.

( $\widehat{H}_1$ ) There exist  $M_1, M_3 > 0$ , a continuous function  $u: I \rightarrow \mathbb{R}^+$  and a nondecreasing function  $L: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \leq u(t)L(\|y\|_J), \quad \text{for } t \in I, y \in C^+(J)$$

and also

$$\phi(0) + L(M_i) \left( \zeta r_i \int_0^1 (Hu)(s) dg(s) + \int_0^{r_i} (Hu)(s) ds \right) < M_i r_i, \quad i = 1, 3.$$

( $\widehat{H}_2$ ) There exist a constant  $\delta \in (0, 1)$  and functions  $\tau: I \rightarrow [0, q]$ , continuous  $v: I \rightarrow \mathbb{R}^+$  and nondecreasing  $w: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \geq v(t)w(y(-\tau(t))), \quad t \in X, y \in \{h \in C^+(J) : \|h\|_J < \min\{M_1, M_3\}\},$$

where  $X := \{t \in I : \delta \leq t - \tau(t) \leq 1\}$  and  $M_1, M_3$  are defined in ( $\widehat{H}_1$ ).

( $\widehat{H}_3$ ) There exists  $\rho > 0$  such that

$$\frac{\rho}{\delta} < \phi(0) + w(\rho) \left( r_2 \zeta \int_0^1 (H_X v)(s) dg(s) + \int_0^{r_2} (H_X v)(s) ds \right).$$

Notice that if  $\phi(0) \neq 0$ , then this  $\rho$  always exists.

Using the assumptions ( $\widehat{H}_1$ )–( $\widehat{H}_3$ ) and, in the same way as in the above Theorem 3.2, we can prove the following theorem.

**THEOREM 3.3.** *Suppose that assumptions ( $\widehat{H}_1$ )–( $\widehat{H}_3$ ) hold and furthermore  $\|\phi\|_J \leq \min\{M_1, M_3\}$ . Then the boundary value problem (1.1)–(1.3) has at least two concave and nondecreasing on  $I$  and positive on  $J \cup I$  solutions  $x_1, x_2$  such that  $x_1(r_3) > M_3 r_3$ ,  $x_1(r_2) < \rho/\delta$ ,  $x_2(r_2) > \rho/\delta$  and  $x_2(r_1) < M_1 r_1$ .*

The obtained solutions  $x_1, x_2$  in Theorems 3.2 and 3.3 are all nondecreasing. Thus, in the special case when  $r_1 = r_2 = r_3 = 1$ , we have that  $x_i(r_j) = x_i(1) = \|x_i\|$ ,  $i = 1, 2, j = 1, 2, 3$ . Therefore, we have the following corollary of Theorems 3.2 and 3.3.

**COROLLARY 3.4.** *Suppose that assumptions ( $H_1$ )–( $H_3$ ) (resp. ( $\widehat{H}_1$ )–( $\widehat{H}_3$ )) hold and furthermore  $\|\phi\|_J \leq M$  (resp.  $\|\phi\|_J \leq \min\{M_1, M_3\}$ ). Then the boundary value problem (1.1)–(1.3) has at least two concave and nondecreasing on  $I$  and positive on  $J \cup I$  solutions  $x_1, x_2$  such that*

$$\frac{\rho_3}{\delta} < \|x_1\| < M < \|x_2\| < \frac{\rho_1}{\delta} \quad \left( \text{resp. } M_3 < \|x_1\| < \frac{\rho}{\delta} < \|x_2\| < M_1 \right).$$

It is remarkable to observe that this corollary can also be obtained by applying twice the Krasnoselskii's theorem under the same assumptions  $(H_1)$ – $(H_3)$  (resp.  $(\widehat{H}_1)$ – $(\widehat{H}_3)$ ).

#### 4. The ordinary differential equations case

In this section we suppose that  $q = 0$ . Then  $J = \{0\}$ , so the boundary value problem (1.1)–(1.3) is reformulated as follows

$$(4.1) \quad x''(t) + f(t, x(t)) = 0 \quad \text{for } t \in I,$$

$$(4.2) \quad x(0) = N,$$

$$(4.3) \quad x'(1) = \int_0^1 x'(s) dg(s),$$

where  $f: \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a continuous function,  $g: I \rightarrow \mathbb{R}^+$  is a nondecreasing function, such that  $g(0) = 0$ ,  $1 - g(1) > 0$  and  $N \in \mathbb{R}^+$ . Note that equation (4.1) is equivalent to the following form

$$x''(t) + f(t, x_t(0)) = 0 \quad \text{for } t \in I$$

and  $C^+(\{0\}) \equiv \mathbb{R}^+$ , so  $f: \mathbb{R}^+ \times C^+(\{0\}) \rightarrow \mathbb{R}^+$ .

Now, the analogue of Lemma 2.2 for this case is the following:

LEMMA 4.1. *A function  $x \in C(I)$  is a solution of the boundary value problem (4.1)–(4.3) if and only if  $x$  is a fixed point of the operator  $\widehat{A}: C(I) \rightarrow C(I)$ , with*

$$\widehat{A}x(t) = N + \zeta t \int_0^1 \int_s^1 f(r, x(r)) dr dg(s) + \int_0^t \int_s^1 f(r, x(r)) dr ds, \quad t \in I,$$

where  $\zeta := 1/(1 - g(1))$ .

Assumptions  $(H_1)$ – $(H_3)$  and  $(\widehat{H}_1)$ – $(\widehat{H}_3)$ , for the special case  $q = 0$ , are stated as follows:

$(H_1)_0$  There exist  $M > 0$ , continuous function  $u: I \rightarrow \mathbb{R}^+$  and nondecreasing function  $L: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \leq u(t)L(y) \quad \text{for } t \in I, y \in \mathbb{R}^+$$

and

$$N + L(M) \left( \zeta r_2 \int_0^1 (Hu)(s) dg(s) + \int_0^{r_2} (Hu)(s) ds \right) \leq Mr_2,$$

where the function  $H$  is defined in the previous section.

$(H_2)_0$  There exist  $\delta \in (0, 1)$  and functions  $v: I \rightarrow \mathbb{R}^+$  continuous and  $w: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  nondecreasing such that

$$f(t, y) \geq v(t)w(y) \quad \text{for } t \in Z := [\delta, 1], y \in [0, M].$$

(H<sub>3</sub>)<sub>0</sub> There exist  $\rho_1, \rho_3 > 0$  such that

$$\frac{\rho_i}{\delta} \leq N + w(\rho_i) \left( r_i \zeta \int_0^1 (H_Z v)(s) dg(s) + \int_0^{r_i} (H_Z v)(s) ds \right), \quad i = 1, 3,$$

where the function  $H_Z$  is defined in previous section.

( $\widehat{H}_1$ )<sub>0</sub> There exist  $M_1, M_3 > 0$ , a continuous function  $u: I \rightarrow \mathbb{R}^+$  and a nondecreasing function  $L: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \leq u(t)L(y) \quad \text{for } t \in I, y \in \mathbb{R}^+$$

and

$$N + L(M_i) \left( \zeta r_i \int_0^1 (Hu)(s) dg(s) + \int_0^{r_i} (Hu)(s) ds \right) < M_i r_i, \quad i = 1, 3$$

where the function  $H$  is defined in the previous section.

( $\widehat{H}_2$ )<sub>0</sub> There exist a constant  $\delta \in (0, 1)$ , a continuous function  $v: I \rightarrow \mathbb{R}^+$  and a nondecreasing function  $w: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$f(t, y) \geq v(t)w(y) \quad \text{for } t \in Z := [\delta, 1], y \in \mathbb{R}^+.$$

( $\widehat{H}_3$ )<sub>0</sub> There exists  $\rho > 0$  such that

$$\frac{\rho}{\delta} < N + w(\rho) \left( r_2 \zeta \int_0^1 (H_Z v)(s) dg(s) + \int_0^{r_2} (H_Z v)(s) ds \right),$$

where the function  $H_Z$  is defined in the previous section.

Therefore, we have the following theorems, which are analogues of Theorems 3.2 and 3.3, respectively.

**THEOREM 4.2.** *Suppose that assumptions ( $\widehat{H}_1$ )–( $\widehat{H}_3$ ) hold and furthermore  $N \leq M$ . Then the boundary value problem (4.1)–(4.3) has at least two concave, nondecreasing and positive on  $I$  solutions  $x_1, x_2$  such that  $x_1(r_3) > \rho_3/\delta$ ,  $x_1(r_2) < Mr_2$ ,  $x_2(r_2) > Mr_2$  and  $x_2(r_1) < \rho_1/\delta$ .*

**THEOREM 4.3.** *Suppose that assumptions ( $\widehat{H}_1$ )<sub>0</sub>–( $\widehat{H}_3$ )<sub>0</sub> hold and furthermore  $N \leq \min\{M_1, M_3\}$ . Then the boundary value problem (4.1)–(4.3) has at least two concave, nondecreasing and positive on  $I$  solutions  $x_1, x_2$  such that  $x_1(r_3) > M_3 r_3$ ,  $x_1(r_2) < \rho/\delta$ ,  $x_2(r_2) > \rho/\delta$  and  $x_2(r_1) < M_1 r_1$ .*

Also, the following corollary corresponds to Corollary 3.4.

**COROLLARY 4.4.** *Suppose that assumptions (H<sub>1</sub>)<sub>0</sub>–(H<sub>3</sub>)<sub>0</sub> (resp. ( $\widehat{H}_1$ )<sub>0</sub>–( $\widehat{H}_3$ )<sub>0</sub>) hold and furthermore  $N \leq M$  (resp.  $N \leq \min\{M_1, M_3\}$ ). Then the boundary value problem (4.1)–(4.3) has at least two concave, nondecreasing and positive on  $I$  solutions  $x_1, x_2$  such that*

$$\frac{\rho_3}{\delta} < \|x_1\| < M < \|x_2\| < \frac{\rho_1}{\delta} \quad \left( \text{resp. } M_3 < \|x_1\| < \frac{\rho}{\delta} < \|x_2\| < M_1 \right).$$

## 5. Applications

5.1. Consider the boundary value problem

$$(5.1) \quad x''(t) + \left[ x\left(t - \frac{1}{2}\right) - \frac{4}{5} \right]^5 + 1 = 0 \quad \text{for } t \in I := [0, 1],$$

$$(5.2) \quad x_0(t) = \phi(t) := t^2 \quad \text{for } t \in J := \left[-\frac{1}{2}, 0\right],$$

$$(5.3) \quad x'(1) = \int_0^1 x'(s) dg(s),$$

where

$$g(t) = \begin{cases} 0 & \text{for } 0 \leq t \leq \frac{1}{3}, \\ \frac{3}{2}t - \frac{1}{2} & \text{for } \frac{1}{3} \leq t \leq \frac{1}{2}, \\ \frac{1}{2}t & \text{for } \frac{1}{2} \leq t \leq 1. \end{cases}$$

It is easy to see that condition (5.3) is equivalent to the following nonlocal one

$$x'(1) = \frac{1}{2}x(1) + x\left(\frac{1}{2}\right) - \frac{3}{2}x\left(\frac{1}{3}\right).$$

Obviously,  $f(t, y) := (y - 4/5)^5 + 1$  is nonnegative on  $\mathbb{R}^+ \times C^+(J)$ ,  $\phi$  is nonnegative on  $J$  and  $g$  is nondecreasing, with  $g(0) = 0$  and  $1 - g(1) = 1/2 > 0$ . Set  $r_1 = 2/5$ ,  $r_2 = 3/5$  and  $r_3 = 4/5$ . Define  $L(z) = (z - 4/5)^5 + 1$ ,  $z \in \mathbb{R}^+$ , and  $u(t) = 1$ ,  $t \in I$ . Since inequality

$$L(M) < \frac{60}{67}M$$

holds for  $M = 7/5$ , assumption (H<sub>1</sub>) is satisfied.

Additionally, set  $\delta = 1/4$ ,  $\tau(t) = 1/2$ ,  $t \in I$ ,  $v(t) = 1$ ,  $t \in I$  and  $w(z) = (z - 4/5)^5 + 1$ ,  $z \in \mathbb{R}^+$ . Then,  $X = [3/4, 1]$  and the inequalities in (H<sub>3</sub>) assume, take the form

$$w(\rho_1) > \frac{64}{3}\rho_1 \quad \text{and} \quad w(\rho_3) > \frac{3200}{299}\rho_3,$$

which are satisfied for  $\rho_1 = 6$  and  $\rho_3 = 1/20$ . Finally, it is obvious that  $\|\phi\|_J \leq 7/5$ , so we can apply Theorem 3.2 to get that the boundary value problem (5.1)–(5.3) has at least two concave and nondecreasing on  $I$  and positive on  $J \cup I$  solutions  $x_1, x_2$ , such that

$$x_1\left(\frac{4}{5}\right) > \frac{1}{5}, \quad x_1\left(\frac{3}{5}\right) < \frac{21}{25}, \quad x_2\left(\frac{3}{5}\right) > \frac{21}{25} \quad \text{and} \quad x_2\left(\frac{2}{5}\right) < 24.$$

**5.2.** Once again, consider the boundary value problem (5.1)–(5.3).

Having in mind Corollary 3.4, we set  $r_1 = r_2 = r_3 = 1$ . Define also again  $L(z) = (z - 4/5)^5 + 1$ ,  $z \in \mathbb{R}^+$ , and  $u(t) = 1$ ,  $t \in I$ . Since inequality

$$L(M) < \frac{12}{11}M$$

holds for  $M = 11/10$ , assumption  $(H_1)$  is satisfied.

Additionally, set  $\delta = 1/4$ ,  $\tau(t) = 1/2$ ,  $t \in I$ ,  $v(t) = 1$ ,  $t \in I$  and  $w(z) = (z - 4/5)^5 + 1$ ,  $z \in \mathbb{R}^+$ . Then,  $X = [3/4, 1]$  and the inequalities in assumption  $(H_3)$  take the forms

$$w(\rho_1) > \frac{64}{7}\rho_1 \quad \text{and} \quad w(\rho_3) > \frac{64}{7}\rho_3,$$

which are satisfied for  $\rho_1 = 27/10$  and  $\rho_3 = 1/12$ . Finally, it is obvious that  $\|\phi\|_J \leq 11/10$ , so we can apply Corollary 3.4 to get that the boundary value problem (5.1)–(5.3) has at least two concave and nondecreasing on  $I$  and positive on  $J \cup I$  solutions  $x_1, x_2$ , such that

$$\frac{1}{3} < \|x_1\| < \frac{11}{10} < \|x_2\| < \frac{54}{5}.$$

**5.3.** Consider the boundary value problem

$$(5.4) \quad x''(t) + 8 \arctan(10x(t) - 14) + 12 = 0 \quad \text{for } t \in I := [0, 1],$$

$$(5.5) \quad x(0) = 0,$$

$$(5.6) \quad x'(1) = \int_0^1 x'(s) dg(s),$$

where  $g(t) = t/4$ ,  $t \in I$ .

It is clear that (5.6) is equivalent to the boundary condition

$$x'(1) = \frac{1}{4} \left( x(1) - x(0) \right).$$

Obviously,  $f(t, y) := 8 \arctan(10y - 14) + 12$  is positive on  $\mathbb{R}^+ \times \mathbb{R}^+$  and  $g$  is nondecreasing, with  $g(0) = 0$  and  $1 - g(1) = 3/4 > 0$ . Set  $r_1 = 1/4$ ,  $r_2 = 1/2$  and  $r_3 = 3/4$ . Define  $L(z) = 8 \arctan(10z - 14) + 12$ ,  $z \in \mathbb{R}^+$ , and  $u(t) = 1$ ,  $t \in I$ . Since inequalities

$$L(M_1) < \frac{24}{25}M_1 \quad \text{and} \quad L(M_3) < \frac{24}{19}M_3$$

hold for  $M_1 = 30$  and  $M_3 = 1/100$ , assumption  $(\widehat{H}_1)_0$  is satisfied.

Additionally, set  $\delta = 1/2$ ,  $v(t) = 1$ ,  $t \in I$  and  $w(z) = 8 \arctan(10z - 14) + 12$ ,  $z \in \mathbb{R}^+$ . Then,  $Z = [1/2, 1]$  and assumption  $(\widehat{H}_3)_0$  takes the form

$$w(\rho) > \frac{96}{15}\rho,$$

which is satisfied for  $\rho = 7/5$ . Finally, it is obvious that  $N = 0 \leq 1/100 = \min\{M_1, M_3\}$ , so we can apply Theorem 4.3 to get that the boundary value problem (5.4)–(5.6) has at least two concave and nondecreasing and positive on  $I$  solutions  $x_1, x_2$ , such that

$$x_1\left(\frac{3}{4}\right) > \frac{3}{400}, \quad x_1\left(\frac{1}{2}\right) < \frac{14}{5}, \quad x_2\left(\frac{1}{2}\right) > \frac{14}{5} \quad \text{and} \quad x_2\left(\frac{1}{4}\right) < \frac{15}{2}.$$

## REFERENCES

- [1] R. I. AVERY AND J. HENDERSON, *Two positive fixed points of nonlinear operators on ordered Banach spaces*, Comm. Appl. Nonlinear Anal. **8** (2001), 27–36.
- [2] R. I. AVERY, CHUAN JEN CHYAN AND J. HENDERSON, *Twin solutions of boundary value problems for ordinary differential equations and finite difference equations*, Comput. Math. Appl. **42** (2001), 695–704.
- [3] N. AZBELEV, V. MAKSIMOV AND L. RAKHMATULLINA, *Introduction to the Theory of Linear Functional Differential Equation*, World Federation Publishers Co., Atlanta, Georgia, 1995.
- [4] J. K. HALE AND S. M. V. LUNEL, *Introduction to Functional Differential Equations*, Springer–Verlag, New York, 1993.
- [5] J. HENDERSON, *Double solutions of three-point boundary-value problems for second-order differential equations*, Electron. J. Differential Equations. **2004** (2004), no. 115, 1–7.
- [6] G. L. KARAKOSTAS, K. G. MAVRIDIS AND P. CH. TSAMATOS, *Multiple positive solutions for a functional second-order boundary-value problem*, J. Math. Anal. Appl. **282** (2003), 567–577.
- [7] ———, *Triple solutions for a nonlocal functional boundary-value problem by Leggett–Williams theorem*, Appl. Anal. **83** (2004), 957–970.
- [8] G. L. KARAKOSTAS AND P. CH. TSAMATOS, *Positive solutions for a nonlocal boundary-value problem with increasing response*, Electron. J. Differential Equations **2000** (2000), no. 73, 1–8.
- [9] ———, *Multiple positive solutions for a nonlocal boundary-value problem with response function quiet at zero*, Electron. J. Differential Equations **2001** (2001), no. 13, 1–10.
- [10] ———, *Sufficient conditions for the existence of nonnegative solutions of a nonlocal boundary value problem*, Appl. Math. Lett. **15** (2002), 401–407.
- [11] ———, *Uniformly quiet at zero functions and existence results for one-parameter boundary value problems*, Ann. Polon. Math. **78** (2002), 267–276.
- [12] ———, *Positive solutions and nonlinear eigenvalue problems for retarded second order differential equations*, Electron. J. Differential Equations **2002** (2002), no. 59, 1–11.
- [13] M. A. KRASNOSEL'SKIĬ, *Positive Solutions of Operator Equations*, Noordhoff, Groningen, 1964.
- [14] R. W. LEGGETT AND L. R. WILLIAMS, *Existence of multiple positive fixed points of nonlinear operators in ordered Banach spaces*, Indiana Univ. Math. J. **28** (1979), 673–688.
- [15] YONGKUN LI AND LIFEI ZHU, *Positive periodic solutions of nonlinear functional differential equations*, Appl. Math. Comp. **156** (2004), 329–339.
- [16] PING LIU AND YONGKUN LI, *Multiple positive periodic solutions of nonlinear functional differential system with feedback control*, J. Math. Anal. Appl. **288** (2003), 819–832.

- [17] PING LIU, YONGKUN LI AND LIGHONG LU, *Positive solutions of nonlocal boundary value problem for nonlinear retarded differential equation*, Appl. Math. J. Chinese Univ. Ser.B **19** (2004), 263–271.
- [18] YUJI LIU AND WEIGAO GE, *Double positive solutions of fourth-order nonlinear boundary value problems*, Appl. Anal. **82** (2003), 369–380.
- [19] ———, *Twin positive solutions of three-point boundary value problems for finite difference equations*, Soochow J. Math. **30** (2004), 11–19.
- [20] P. CH. TSAMATOS, *Positive solutions with given slope of a nonlocal second order boundary value problem with sign changing nonlinearities*, Ann. Pol. Math. **83** (2004), 231–242.
- [21] J. WANG, *The existence of positive solutions for one-dimensional  $p$ -Laplacian*, Proc. Amer. Math. Soc. **125** (1997), 2275–2283.
- [22] E. ZEIDLER, *Nonlinear Functional Analysis and Its Applications I: Fixed-Point Theorems*, Springer–Verlag, New York, 1993.

*Manuscript received June 17, 2005*

PANAGIOTIS CH. TSAMATOS  
Department of Mathematics  
University of Ioannina  
451 10 Ioannina, GREECE  
*E-mail address:* ptsamato@cc.uoi.gr